



An Accurate Taylors Series Solution with High Radius of Convergence for the Blasius Function and Parameters of Asymptotic Variation

S.Anil Lal¹ and Neeraj Paul M²

Department of Mechanical Engineering,
College of Engineering Trivandrum, Kerala, India

† Corresponding Author Email: anillal65@hotmail.com

(Received July, 18, 2013; accepted November, 08, 2013)

ABSTRACT

This paper reports a high accurate solution of the Blasius function $f(\eta)$ in the form of a converging Taylor's series for a higher range of $\eta \in [0, 9]$. The method used consists of conversion of the boundary value problem into an initial value problem and solution by differential transform method. The initial value of the second derivative of the Blasius function is determined from the final value of first derivative of another function. The final value of first derivative of the latter function is determined by the Taylor's series expansions with center at $\eta = 15$. The series expansion for the Blasius function is obtained with center of expansion at $\eta = 4$, is alternating and is accurately converging for higher values of η , with the number of used for summation equal to 2000. The present expansion is obtained without resorting to approximations and has a higher radius of convergence. The first 200 coefficients of the series, the second derivative of the function at $\eta = 0$, the parameters of the asymptotic solution are reported with 21 decimal places accuracy. The level of accuracy of the results presented is higher than any other results reported so far. This note also reports the mathematical steps involved in the derivation of the similarity variable of Blasius problem.

Keywords: Differential transform, Taylor's series, Similarity variable, Blasius function, Exact solution

1. INTRODUCTION

Ever since the formulation and solution of the function defined by $f'''(\eta) + \frac{1}{2}f(\eta)f''(\eta) = 0$, with $f(0) = f'(0) = 0$, $f'(\infty) = 1$ and $\eta \in [0, \infty]$ by Blasius (1908), more than a century ago, the function called as Blasius function has been a subject of great interest to Mathematicians and Physicists (K.Toepfer and H.Blasius. (1912), Howarth (1938), Cortell (2005), Boyd (2008), Zaimi, Bidin, Bakar, and Hamid (2012)), to name a few. These investigations considered analytical as well as numerical methods for the solution of $f(\eta)$. Though the problem has been approximately solved using numerical techniques, difficulties encounter when the exact solution of the function is sought. This is mainly due to the fact that the exact solution is a power series without having a closed form and singularity of the function. A systematic investigation of the simple pole of the function and convergence behavior is reported by Boyd (1999). While a certain minimum number of terms are required to be summed for getting the results with required accu-




Fig. 1. Boundary Layer over a Flat plate.

racy levels, the solution for $\eta \gg 1$ is subjected to convergence problems. The present work explores the potential of differential transform method for solving non-linear initial value problems of functions with singularity. In order to obtain a higher radius of convergence of the series expansion of functions with singularity, shifting of the center of expansion is proposed. A new methodology consisting of recursive application of differential transform method for shifting the center of the expansion has been demonstrated. Hence a high accurate solution of Blasius function in the form of a con-

verging power series with center of expansion at $\eta = 4$ and a higher radius of convergence $\eta \in [0, 9]$ than other results reported so far is obtained. The important parameters of the asymptotic solution are reported with a higher level of accuracy. This technical brief reports the method for converting the boundary value problem into an initial value problem. The first part of this note describes a systematic method for deriving the similarity variable for transforming the PDE of boundary layer equation into the Blasius equation.

2. BOUNDARY LAYER OVER A FLAT PLATE

Figure 1 shows a schematic diagram of boundary layer over a flat plate. It is a zero pressure gradient flow as the potential flow velocity is a constant and equal to the free stream velocity U_∞ . Thus the governing partial differential equation of boundary layer flow over a flat plate is written by neglecting the pressure gradient term in the Prandtl boundary layer equation as:

$$\text{continuity : } \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \tag{1}$$

$$\text{x-momentum : } u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = \nu \frac{\partial^2 u}{\partial y^2} \tag{2}$$

The boundary conditions are $u = v = \frac{\partial^2 u}{\partial y^2} = 0$ at $y = 0$ and $\frac{\partial u}{\partial y} = 0$ at $y = \infty$.

The velocity u in the boundary layer is a function of x and y coordinates. Typical velocity profiles as a function of y drawn for different values of x are shown in Fig. 2(a). A dimensionless variable η formed by combining the independent variables x, y and the physical constants of the problem such that variation of the non-dimensional velocity u/U_∞ as a function of η is a single curve as shown in Fig. 2(b) is called a *similarity variable*. Blasius (1908) defined the similarity variable as $\eta = y/\delta$ using physical arguments based on momentum transport. Cengel (2006) noted that finding of such a variable, assuming it exists, is more of an art than science, and it requires to have a good insight of the problem. But from a mathematical point of view, similarity variable can be visualized as a change of variable for a possible transformation of a PDE into an ODE. The first part of this note illustrates a systematic method for deriving the similarity variable for flat plate boundary layer from a mathematical point of view. A fundamental choice for combining x, y and the physical constants may be to write $\eta = kA(x)B(y)$, where k is related to the physical




Fig. 2. Boundary Layer.

constants such as ν and U_∞ . By noting that all boundary conditions of the problem are based on the values of y , a convenient choice for $B(y)$ is taken as the variable y itself. Thus,

$$\eta = kA(x)y \quad \text{with, } \frac{\partial \eta}{\partial x} = kyA' \quad \text{and} \quad \frac{\partial \eta}{\partial y} = kA \tag{3}$$

The equation of the dimensionless velocity as shown in Fig. 2(b) has the form

$$\frac{u}{U_\infty} = f'(\eta) \quad \text{or} \quad u = U_\infty f'(\eta) \tag{4}$$

It is possible to satisfy the continuity Eq. (1) by deriving the velocity components from stream function $\psi(x, y)$ as

$$u = U_\infty f'(\eta) = \frac{\partial \psi}{\partial y} \Rightarrow \psi = \frac{U_\infty f}{kA} \tag{5}$$

$$v = -\frac{\partial \psi}{\partial x} = -\frac{U_\infty}{k} \left[\frac{A f' k A' y - f A''}{A^2} \right] \tag{6}$$

The other derivatives u are

$$\frac{\partial u}{\partial x} = U_\infty f'' k A' y, \quad \frac{\partial u}{\partial y} = U_\infty f'' k A$$

and

$$\frac{\partial^2 u}{\partial y^2} = U_\infty k^2 A^2 f''' \tag{7}$$

Putting Eq. (5-7) in Eq. (2), it is possible to write

$$f''' - \frac{U_\infty A'}{vk^2 A^3} f f'' = 0 \tag{8}$$

The condition for the above differential equation to represent an ODE is that, $\frac{U_\infty A'}{vk^2 A^3}$ is a constant.

Taking $\frac{U_\infty A'}{vk^2 A^3} = -p$, (p is a constant)

$$\Rightarrow A^{-3} \frac{dA}{dx} = -\frac{pvk^2}{U_\infty}$$

(On integration) $\Rightarrow \frac{1}{A^2} = \frac{2pvk^2 x}{U_\infty}$

$$\Rightarrow A^2 k^2 = \frac{U_\infty}{2pvx} \tag{9}$$

Since p can have any arbitrary constant value, a choice that turns the expression of Ak to a compact form is $p = 1/2$. Thus, the final expressions for Ak and η become

$$kA(x) = \sqrt{\frac{U_\infty}{vx}} \quad \text{and} \quad \eta = y\sqrt{\frac{U_\infty}{vx}} \tag{10}$$

The resulting ODE from Eq. (8) is called the Blasius equation

$$f''' + \frac{1}{2} f f'' = 0; \quad f(0) = f'(0) = 0 \quad \text{and} \quad f'(\infty) = 1 \tag{11}$$

Presented in this section is a straight forward mathematical method for the derivation of similarity variable, for converting a PDE into an ODE. This method is based on mathematical concepts and the application of physics, intuitive judgments and experience are minimum. Existence or suitability of a similarity variable can be judged during the process of transformation.

The following sections consider conversion of the above boundary value problem into an initial value problem and subsequently solution by considering the singularity point of the function, $f(\eta)$. The potential of differential transform Zhou (1986) method is explored to derive accurate solution of the function with a higher radius of convergence.

3. CONVERTING BOUNDARY VALUE PROBLEM (BVP) TO INITIAL VALUE PROBLEM (IVP)

Blasius problem is a non-linear, third order BVP. Solution of a non-linear BVP is complex, either involving iterative techniques or system of equations

along with the use of a numerical technique. At the same time, Initial Value Problems can be solved easily. An important property of Blasius equation (11) is that it is satisfied by $af(a\eta)$ also, which is a scaled variant of $f(\eta)$, where a is a scaling constant. This property called dilational or scaled symmetry of Blasius equation is first observed by Toepfer[2], and this property of the function can be applied to determine the value of $f''(0)$ from the known condition $f'(\infty) = 1$. By knowing the value of $f''(0)$, Blasius problem can be solved as an IVP. The derivation of an expression for $f''(0)$ is reviewed here by considering a function g with

$$g''' + \frac{1}{2} g g'' = 0, \quad \text{with} \\ g(0) = g'(0) = 0 \quad \text{and} \quad g''(0) = 1 \tag{12}$$

By scaled symmetry, the solution of g and f are related by the scaling relation

$f(\eta) = ag(a\eta)$ By differentiating,

we can write $f'(\eta) = a^2 g'(a\eta)$

$$\Rightarrow f'(\infty) = a^2 g'(\infty)$$

$$\Rightarrow g'(\infty) = \frac{f'(\infty)}{a^2} = \frac{1}{a^2}$$

$$\Rightarrow a = [g'(\infty)]^{-1/2}$$

$$f''(\eta) = a^3 g''(a\eta)$$

$$\Rightarrow f''(0) = a^3 g''(0) = a^3 = [g'(\infty)]^{-3/2} \tag{13}$$

The value of $g'(\infty)$ is to be first determined by solving the IVP of g in Eq. (12). By finding $f''(0) = [g'(\infty)]^{-3/2}$, the Blasius problem can be solved as an IVP.

3.1. Solution of IVP

Initial value problems can be solved either numerically or analytically. Blasius IVP has been numerically solved by applying Runge-Kutta method in (K.Toepfer and H.Blasius. (1912), Howarth (1938), Cortell (2005), Boyd (2008), Zaimi, Bidin, Bakar, and Hamid (2012)). Taylor's series method has been used to analytically solve different types of IVPs in (Ganji, Babazadeh, Noori, Pirouz, and Janipour (2009), Asaithambi (2005)). In Taylor's series method, the differential equation is successively differentiated to find higher order derivatives from the known values of lower order derivatives. The series may be a terminating or a non-terminating one, depending on whether the derivatives vanish or not beyond a particular order derivative. A difficulty with non-terminating series is

from its divergence behavior, when the function contains singularities. The radius of convergence of series expansion of a function with singularities is the distance between the center of expansion and its closest singularity. It is possible to change the radius of convergence by changing the center of expansion. A systematic approach for obtaining the Taylor's series solution of IVPs, is first developed by Zhou (1986), called differential transform method. A review of this method and its application for determining the value of $f''(0)$ with the help of the solution of IVP in Eq.(12) are given in the following.

3.1.1. Method of differential transform

The Taylor's expansion of a function $g(\eta)$ about $\eta = \eta_i$ is

$$g(\eta) = g(\eta_i) + \sum_{k=1}^{\infty} \frac{1}{k!} [g^k(\eta)]_{\eta=\eta_i} (\eta - \eta_i)^k$$

$$= G(0) + \sum_{k=1}^{\infty} G(k)(\eta - \eta_i)^k \tag{14}$$

Where $G(k) = \frac{1}{k!} [g^k(\eta)]_{\eta=\eta_i}$ for $k > 0$. $G(k)$ denotes the coefficient of k^{th} degree term of Taylor's series expansion of $g(\eta)$ and $G(0) = g(\eta_i)$. Zhou (1986) considered $G(k)$ as a transform of $g(\eta)$ called *differential transform*. Since then several useful properties of the differential transform were studied (Arikoglu and Ozkol (2005), Bildik and Konuralp (2011), Arikoglu and Ozkol (2008)). These properties have been useful for the solution of IVPs. The differential transform of a derivative $g'(\eta)$ is obtained by differentiating Eq. (14)

$$g'(\eta) = \sum_{k=1}^{\infty} G(k)k(\eta - \eta_i)^{k-1}$$

taking $p = k - 1$

$$= \sum_{p=0}^{\infty} G(p+1)(p+1)(\eta - \eta_i)^p$$

$$= \sum_{k=0}^{\infty} G(k+1)(k+1)(\eta - \eta_i)^k$$

$$= \sum_{k=0}^{\infty} G_1(k)(\eta - \eta_i)^k \tag{15}$$

where $G_1(k) = (k+1)G(k+1)$ is the differential transform of the first derivative of $g(\eta)$. In this manner the differential transform of the m^{th} derivative of $g(\eta)$ is

$$G_m(k) = (k+1)(k+2)...(k+m)G(k+m)$$

$$\Rightarrow G(k+m) = \frac{k!}{(k+m)!} G_m(k) \tag{16}$$

This gives a recurrence relation for the successive differential transforms of $g(\eta)$.

The differential transform of the product term in the governing Eq. (12), viz. $g(\eta)g''(\eta)$ is denoted as $GG_2(k)$ and is obtained using Leibnitz rule as in the following.

$$GG_2(k) = \frac{1}{k!} \frac{d^k}{d\eta^k} [g(\eta)g''(\eta)]$$

Expanding n^{th} differential of product using Leibnitz rule,

$$GG_2(k) = \frac{1}{k!} \sum_{i=0}^k \binom{k}{i} \frac{d^i g(\eta)}{d\eta^i} \frac{d^{k-i} g''(\eta)}{d\eta^{k-i}}$$

$$= \frac{1}{k!} \sum_{i=0}^k \binom{k}{i} i! G(i)(k-i)! G_2(k-i)$$

$$\Rightarrow GG_2(k) = \sum_{i=0}^k G(i)G_2(k-i) \tag{17}$$

By knowing $G(0) = 0$, $G'(0) = 0$ and $G''(0) = 1$, the recurrence relations for computing the differential transform of $g(\eta)$ for any value of k are summarized as

$$G(k+1) = \frac{G_1(k)}{k+1}; \quad G_1(k+1) = \frac{G_2(k)}{k+1} \text{ and}$$

$$G_2(k+1) = -\frac{1}{2(k+1)} \sum_{i=0}^k G(k)G_2(k-i) \tag{18}$$

$G(k)$ is computed up to the required number of terms (k) of the series. By knowing $G(k)$, the power series for $g(\eta)$ is obtained as in Eq. (14)

3.1.2. Determination of initial value of $f''(\eta)$

Firstly, we studied the convergence property of the series obtained for $g(\eta)$ by keeping k arbitrarily equal to 1000. It is found that the series of $g(\eta)$ expanded with $\eta_i = 0$ as the center of the expansion starts to diverge for $\eta > 3.8$. So it is inferred that the function $g(\eta)$ has singularity at $\eta < -3.8$. Due to this singularity, we have restricted the computation of $g(\eta)$ using this series for $\Delta\eta < 3.8$. By treating the value of $g(\eta)$, $g'(\eta)$ and $g''(\eta)$ obtained from this series at $\eta = \Delta\eta$ as the initial value, a second series is computed for $\Delta\eta < \eta < 2\Delta\eta$. In this manner a required number of split series have been

computed for equal intervals of η , $\Delta\eta = 1$ up to a maximum value of η , $\eta_m = 15$. The value of $g'(15)$ computed from the last series is taken as $g'(\infty)$ and hence $f''(0) = [g'(\eta_m = 15)]^{-3/2}$ is computed. The result obtained from the present extended double precision computation with 21 decimal digits accuracy and using a total of 40 terms in each series is

$$f''(0) = 0.332057336215196299183$$

The above value is found to be invariant for $\eta_m > 15$ and $k > 40$. The result matches exactly with the value reported by Boyd (1999), which is found numerically using Runge-Kutta method, 27 decimal place accuracy computations and 50,000 grid points. While the computational effort involved in the present solution is much less, it confirms that the results Boyd (1999) believes as correct is correct. However the result obtained in the present work is more accurate than already reported values.

4. SOLUTION OF BLASIUS FUNCTION

With the result for $f''(0)$ obtained above, the Blasius function defined defined by $f''' + \frac{1}{2}ff'' = 0$ with $f(0) = f'(0) = 0$ becomes an initial value problem and is solved by computing $F(k)$ using the recursive relations in Eq.(18) by changing G to F and $F(0) = F_1(0) = 0$ and $F_2(0) = 0.332057336215196299183$. The series thus obtained is the Taylor's series expansion about $\eta = 0$ and is found to diverge for $\eta > 5.69$ due to the singularity of $f(\eta)$ at $\eta = -5.69$, Boyd (1999). Hence the series obtained is used to compute the values of $f(4) = 2.30574641846207118020$, $f'(4) = 0.955518229810694247274$ and $f''(4) = 6.42341210916906681680E - 0002$. Using these values as initial values at $\eta = 4$, another series for $f(\eta)$ is computed with the center of expansion at $\eta = 4$. The series is an alternating one. Computation of the function by summing finite number of terms of the series expanded about $\eta = 4$ shows that, accurate result is obtained in $0 \leq \eta \leq 8$, for summation with only 200 terms. But the results obtained by summing 200 terms of the series is found to give unrealistic values for $\eta > 8$. By increasing the number of terms of summation to 2000, the same series is found to give accurate results up to $\eta = 9$. The values of the function and its derivatives obtained in the present work are tabulated in Table 1. The variation of $f'(\eta)$ and $f''(\eta)$ thus obtained are depicted in Fig. 3. The first 200 coefficients of series are given in Table 2. We expect that the present Taylor series expansion about $\eta = 4$ is more accurate than any other results reported in literature. At $\eta = 9$, the solution is close to asymptotic variation with the value of $f'(\eta)$ being nearly constant with changes only after 7 decimal places.




Fig. 3. Variation of $f'(\eta)$ and $f''(\eta)$ with respect to η .

We also investigated the asymptotic parameters using successive series constructed up to $\eta = 20$ and the asymptotic variation is obtained as

$$f(\eta) = \eta + B \quad \text{and}$$

$$f''(\eta) = Q \exp\left(-\frac{1}{4}\eta(\eta + 2B)\right)$$

where the obtained values of B and Q are -1.72078765752050281274 and 0.1114837549978888824 respectively. The present value of B is found to match with a better accuracy with the results presented in Boyd (2008) of -1.720787657520503 . However, the value of Q obtained is different from the one reported in Boyd (2008) of 0.233727621285063 . The results reported in this paper is after a through verification only.

5. CONCLUSIONS

The method of differential transforms has been applied to generate the Taylor's series expansion of the Blasius function, $f(\eta)$. The paper reports a methodology involving recursive application of differential transform method for shifting the center of expansion of functions expressed in the form of differential equations. The series for Blasius function has been expanded about $\eta = 4$, which provided a higher radius of convergence in the positive range of η . The series is alternating and the accuracy of the evaluated function is found to depend on the number of terms summed for $\eta > 4$. The series reported is convergent for $\eta \in [0, 9]$ when 2000 are used for summation. A 21 decimal places accurate value of the second derivative of the function at $\eta = 0$, which physically represents the shear stress on the surface of the plate is 0.332057336215196299183 . The parameters of the asymptotic solution is also obtained with a 21 decimal places accuracy.

Table 1 Tabulated values of $f(\eta)$, $f'(\eta)$, $f''(\eta)$ obtained with Taylor's series expansion about $\eta = 4$

η	$f(\eta)$	$f'(\eta)$	$f''(\eta)$
0	-2.90451215935441549708E-0019	2.92231574613161685905E-0019	0.332057336215196
0.2	6.64099971459704033365E-0003	6.64077920962507645673E-0002	0.331983837114628
0.4	2.65598840179947234234E-0002	0.132764160761022	0.331469844201455
0.6	5.97346374980383965769E-0002	0.198937252422219	0.330079127574296
0.8	0.106108220839041	0.264709138723117	0.327389270149252
1	0.165571725789280	0.329780031249667	0.323007116686943
1.2	0.237948717288927	0.393776104433956	0.316589191061116
1.4	0.322981573829535	0.456261764705134	0.307865391790168
1.6	0.420320765501626	0.516756784422615	0.296663461455719
1.8	0.529518037743810	0.574758143889457	0.282931017259756
2	0.650024369935289	0.629765736502386	0.266751545697278
2.2	0.781193337010573	0.681310377236027	0.248350913190371
2.4	0.922290125634541	0.728981935062575	0.228091760686684
2.6	1.072505976787830	0.772455021148565	0.206454626799426
2.8	1.230977302314310	0.811509623199109	0.184006593865361
3	1.396808230870350	0.846044443657994	0.161360319540879
3.2	1.569094960006760	0.876081455172477	0.139128055572426
3.4	1.746950093948840	0.901761221436765	0.117876246087524
3.6	1.929525169760530	0.923329665881649	9.80862787842802132048E-0002
3.8	2.116029817172070	0.941117996729313	8.01259181391971851178E-0002
4	2.305746418462070	0.95518229810694	6.42341210916906681680E-0002
4.2	2.498039662706970	0.966957073753606	5.05197474866457608118E-0002
4.4	2.692360937543090	0.975870832136478	3.89726108536295503280E-0002
4.6	2.888247990017810	0.982683500760700	2.94837720116487089967E-0002
4.8	3.085320655177480	0.987789526200787	2.18711863474432831236E-0002
5	3.283273665156310	0.991541900164393	1.59067986853181631358E-0002
5.2	3.481867611508660	0.994245535359928	1.13417889689291312873E-0002
5.4	3.680919062897510	0.996155303962753	7.92765981470613569935E-0003
5.6	3.880290677633380	0.997477768212915	5.43195767992749243316E-0003
5.8	4.079881939239700	0.998375493650191	3.64841366674732585084E-0003
6	4.279620922513850	0.998972872435860	2.40203984375727885487E-0003
6.2	4.479457297282690	0.999362541719091	1.55017069065636743797E-0003
6.4	4.679356615431430	0.999611701717676	9.80615117009763015151E-0004
6.6	4.879295811060250	0.999767870210032	6.08044264784485316850E-0004
6.8	5.079259772449010	0.999863819037038	3.69562570140311736630E-0004
7	5.279238811029110	0.999921604147950	2.20168955271134609014E-0004
7.2	5.479226847343180	0.999955717277927	1.28569807235158993429E-0004
7.4	5.679220147333840	0.999975457684898	7.35929833892270373715E-0005
7.6	5.879216465804060	0.999986655139129	4.12903111137017749811E-0005
7.8	6.079214481071950	0.999992881166101	2.27077514028069458891E-0005
8	6.279213431346080	0.999996274535301	1.22409262432531429552E-0005
8.2	6.479212886678500	0.999998087459238	6.46797861084886973152E-0006
8.4	6.679212609441440	0.999999036874844	3.34993975319999345318E-0006
8.6	6.879212471015070	0.999999524247805	1.70066798857665268096E-0006
8.8	7.079212403216730	0.999999769489751	8.46284121306853119521E-0007
9	7.279212370645330	0.999999890444161	3.92686164651959923121E-0007

Table 2 First 200 coefficients of the Taylor series expansion of the Blasius function given by $f(\eta) = \sum_{k=0} F(k)(\eta - 4)^k$.

k	$F(k)$	k	$F(k)$	k	$F(k)$
0	2.305746418462070	67	1.54501387536272496428E-0047	134	3.32266142258623029852E-0095
1	0.955518229810694	68	-5.31812426339887940710E-0049	135	-1.98310538234261656202E-0095
2	3.21170605458453340840E-0002	69	-5.55103174036789546845E-0049	136	4.91278759876207123111E-0097
3	-1.23422995541853953195E-0002	70	7.08312566937082614087E-0050	137	7.29857631119625434430E-0097
4	2.27859175576908150426E-0003	71	1.52787645522513563236E-0050	138	-8.49971735072191697622E-0098
5	4.70873734551596933229E-0005	72	-4.14304257606461522218E-0051	139	-2.08270073195395656503E-0098
6	-1.04696147203046357531E-0004	73	-2.22689450789485071198E-0052	140	5.19628448478923655496E-0099
7	1.04849728513766856216E-0005	74	1.8180990936048579821E-0052	141	3.44441060727620368746E-0100
8	3.65883591635046807686E-0006	75	-7.70672457610464815904E-0054	142	-2.33913302679774233968E-0100
9	-8.26198757173262535534E-0007	76	-6.40184069216592913088E-0054	143	7.64798370293702712169E-0102
10	-8.56871408032604090498E-0008	77	8.78278560115632077167E-0055	144	8.44177733364167930062E-0102
11	4.24206814475960916804E-0008	78	1.70679514631881930489E-0055	145	-1.05983538572098169671E-0102
12	1.24757464648407639619E-0010	79	-4.96792309934810891449E-0056	146	-2.33973464442283964453E-0103
13	-1.73812386504572966263E-0009	80	-2.18166098174541908883E-0057	147	6.24732251568732334987E-0104
14	1.32651440735246801945E-0010	81	2.13605639450487796055E-0057	148	3.5000357689636777002E-0105
15	5.77723961725846969334E-0011	82	-1.07501564878974550764E-0058	149	-2.75444920540216826285E-0105
16	-9.81264996803191772378E-0012	83	-7.36873048290769883333E-0059	150	1.11759533406814347334E-0106
17	-1.41677737579443119458E-0012	84	1.08427286052096212108E-0059	151	9.74493755794952565242E-0107
18	4.94795732662933281514E-0013	85	1.89852773644347656379E-0060	152	-1.31507152738594442086E-0107
19	1.17217679853551754637E-0014	86	-5.94498871412526580801E-0061	153	-2.61829774025888477805E-0108
20	-1.99587570380654368497E-0014	87	-2.04918333875349427543E-0062	154	7.49489815935327201785E-0109
21	1.32173139876612546061E-0015	88	2.50554860386806934685E-0062	155	3.46247610727717591164E-0110
22	6.50222871563387972142E-0016	89	-1.45989473509745392038E-0063	156	-3.23808598353568341220E-0110
23	-1.09893230989100695430E-0016	90	-8.46414930933756714068E-0064	157	1.56837815247303581412E-0111
24	-1.52737229367497166915E-0017	91	1.33329661960224071757E-0064	158	1.12264456091526251036E-0111
25	5.66376444706610288373E-0018	92	2.10162418626519246400E-0065	159	-1.62464168640150274164E-0112
26	8.23901176700155878733E-0020	93	-7.10027168819215405316E-0066	160	-2.91748745259002246431E-0113
27	-2.28577360586863227790E-0019	94	-1.80033565851515105991E-0067	161	8.97318111790271961126E-0114
28	1.74072742066617737900E-0020	95	2.93411471403056952770E-0067	162	3.29739576663407194300E-0115
29	7.35859232836632572406E-0021	96	-1.94347659185749439227E-0068	163	-3.80028614160509840123E-0115
30	-1.34125625738890900092E-0021	97	-9.70115483508269533097E-0069	164	2.1397871083118307653E-0116
31	-1.66576227379246042319E-0022	98	1.63361932786724523668E-0069	165	1.29058709364809346050E-0116
32	6.72719348653922713935E-0023	99	2.3137279996307639952E-0070	166	-1.99914874450570455454E-0117
33	4.47986953506900797484E-0025	100	-8.46393064540148693651E-0071	167	-3.23538207487299957791E-0118
34	-2.66217397630806611422E-0024	101	-1.39872542971117784716E-0072	168	1.07218295353518230320E-0118
35	2.22094716106009716605E-0025	102	3.43024735807583352017E-0072	169	2.96153117239146438211E-0120
36	8.37518589248022676550E-0026	103	-2.5476257587736166286E-0073	170	-4.45262995301557455429E-0120
37	-1.62032325216119842570E-0026	104	-1.10931882405391600733E-0073	171	2.85915374479122709354E-0121
38	-1.80417114775513806918E-0027	105	1.99498870478017624911E-0074	172	1.48037738674014021808E-0121
39	7.94497605733967138028E-0028	106	2.53129553760214968666E-0075	173	-2.45111629181960645349E-0122
40	-1.31123133608615312516E-0030	107	-1.00708268177728895733E-0075	174	-3.56871470588556566006E-0123
41	-3.08709986811484909863E-0029	108	-7.99094501758040271974E-0078	175	1.27867865023698921137E-0123
42	2.8364351339929736220E-0030	109	4.00347533206144321137E-0077	176	2.4006995010899130421E-0125
43	9.48498873147394786300E-0031	110	-3.29860210791528839734E-0078	177	-5.20816657462216582648E-0125
44	-1.96239700983573131081E-0031	111	-1.26537714511528773244E-0078	178	3.75939661400478264250E-0126
45	-1.93062640742544893606E-0032	112	2.42887538378505860767E-0079	179	1.6941407846743310407E-0126
46	9.39884978039657251875E-0033	113	2.7492650136910772531E-0080	180	-2.99531171891903060726E-0127
47	-9.44658358717036897237E-0035	114	-1.19611735377873318470E-0080	181	-3.91246681071946192982E-0128
48	-3.58235244301780394135E-0034	115	5.45848162805759581119E-0084	182	1.52211748212612817695E-0128
49	3.59998116828591784697E-0035	116	4.6644522909785624503E-0082	183	1.54740743786132968865E-0130
50	1.07319195469366880764E-0035	117	-4.22787751275740452986E-0083	184	-6.08154750822682764854E-0130
51	-2.37289361171275313764E-0036	118	-1.43960886734871669480E-0083	185	4.88007598324053527639E-0131
52	-2.04749652874355638181E-0037	119	2.94875414605123670983E-0084	186	1.93403040466493136928E-0131
53	1.11071547179463322454E-0037	120	2.96063239108923121816E-0085	187	-3.64911916541314266416E-0132
54	-2.02932330331279453972E-0039	121	-1.41812584925099760909E-0085	188	-4.25936402268573012745E-0133
55	-4.15165949145151586119E-0039	122	1.2313897698643687366E-0087	189	1.80862982435781603856E-0133
56	4.53729073624533658245E-0040	123	5.42499798635919130266E-0087	190	3.17975556085456888547E-0136
57	1.21099631862884071177E-0040	124	-5.37309525832258432683E-0088	191	-7.08915855474350595345E-0135
58	-2.86332220243352692683E-0041	125	-1.6332420851599833194E-0088	192	6.26863364229039895060E-0136
59	-2.14341684396764355044E-0042	126	3.57042744191734994042E-0089	193	2.20216391074229466648E-0136
60	1.31092098910088389845E-0042	127	3.15594254328568438522E-0090	194	-4.43296914517726799461E-0137
61	-3.45994909030007761337E-0044	128	-1.67842448710311967717E-0090	195	-4.59925337912476525914E-0138
62	-4.80456683832639116100E-0044	129	2.81616923412489590751E-0092	196	2.14527570427798719383E-0138
63	5.68461628720700586137E-0045	130	6.29823623471267266357E-0092	197	-1.39089427591724692783E-0140
64	1.36252363156128811891E-0045	131	-6.77919326752912501650E-0093	198	-8.24924536692309434289E-0140
65	-3.44793820089044350070E-0046	132	-1.84733958285133870664E-0093	199	7.98184787277333689447E-0141
66	-2.20803805096678201288E-0047	133	4.31239629098361091948E-0094		

REFERENCES

- Arikoglu, A. and I. Ozkol (2005). Solution of boundary value problems for integro-differential equations by using differential transform method. *Applied Mathematics and Computation* 168(2), 1145–1158.
- Arikoglu, A. and I. Ozkol (2008). Solutions of integral and integro-differential equation systems by using differential transform method. *Computers & Mathematics with Applications* 56(9), 2411–2417.
- Asaithambi, A. (2005). Solution of the falkner-skank equation by recursive evaluation of taylor coefficients. *Journal of Computational and Applied Mathematics* 176(1), 203–214.
- Bildik, N. and A. Konuralp (2011). The use of variational iteration method, differential transform method and adomian decomposition method for solving different types of nonlinear partial differential equations. *International Journal of Nonlinear Sciences and Numerical Simulation* 7(1), 65–70.
- Blasius, H. (1908). Grenzschichten in flüssigkeiten mit kleiner reibung. *Z. Math Phys.*, 56:137.
- Boyd, J. (1999). The blasius function in the complex plane. *Experimental Mathematics* 8(4), 381–394.
- Boyd, J. (2008). The blasius function: computations before computers, the value of tricks, undergraduate projects, and open research problems. *SIAM review* 50(4), 791–804.
- Cengel, Y. A. (2006). Heat and mass transfer: A practical approach (si units).
- Cortell, R. (2005). Numerical solutions of the classical blasius flat-plate problem. *Applied mathematics and computation* 170(1), 706–710.
- Ganji, D., H. Babazadeh, F. Noori, M. Pirouz, and M. Janipour (2009). An application of homotopy perturbation method for non-linear blasius equation to boundary layer flow over a flat plate. *International Journal of Nonlinear Science* 7(4), 399–404.
- Howarth, L. (1938). On the solution of the laminar boundary layer equations. *Proc London Math Soc A*, 164:54779.
- K.Toepfer and H.Blasius. (1912). Grenzschichten in flüssigkeiten mit kleiner reibung. *Z. Math Phys.(1912)*.
- Zaimi, W., B. Bidin, N. Bakar, and R. Hamid (2012). Applications of runge-kutta-fehlberg method and shooting technique for solving classical blasius equation. *World Applied Sciences Journal 17 (Special Issue of Applied Math)*, 10–15.
- Zhou, J. (1986). Differential transformation and its applications for electrical circuits.